

IMPACT OF GIBBERELIC ACID TREATMENTS ON GERMINATION OF CLEOPATRA MANDARIN SEEDS

Tarek A. Mahmoud*, Manal A. M. Abo-Eid and Mahmoud A. Khodier

Citrus Department, Horticulture Institute, Agriculture Research Centre, Giza, Egypt.

*Corresponding author: tarekmsc@yahoo.com Received: 26 Aug. 2021 ; Accepted: 25 Sept. 2021

ABSTRACT: This study was carried out throughout two successive seasons (2020 and 2021) for investigating GA₃ pre-sowing soak solutions influence on Cleopatra mandarin seeds (*Citrus reshni* L.). It was aimed to enhance the Cleopatra mandarin seeds germination process, as well as seedlings vigor. Thus, seeds of Cleopatra mandarin rootstock were subjected to soaking for 24 hours in GA₃ solutions at 2000, 2500, and 3000 ppm as well as soaking in tap water (control treatment), for investigating their effect on some germination and growth parameters. Data obtained during both experimental seasons revealed obviously that all GA₃ treatments increased significantly both germination measurements (total germinated seeds & germination %) and some growth parameters as compared to control (soak for 24 hours in tap water). However, soaking in GA₃ at 2500 and 3000 ppm concentrated solutions particularly were statistically the most effective in this study.

Key words: Cleopatra mandarin seeds, gibberellic acid, germination, rootstock, citrus.

INTRODUCTION

Citrus is one of the most important fruit crops all over the world. It ranks the third position between fruit crops and only preceded with grapes and apples. In Egypt, it is the most important fruit crop. Cleopatra mandarin (*Citrus reshni* L.) has been established as rootstock over many years. Tree vigor and fruit quality of sweet orange and mandarin cultivars on Cleopatra mandarin rootstock is very good and its equal to those produced on sour orange rootstock, but fruit size is smaller. Cleopatra mandarin is considered one of the important species of citrus root-stocks, which studies have begun on it as a substitute for the sour orange rootstock in many countries. Cleopatra mandarin is more tolerant than sour orange for Tristeza, gummosis, exocortis, phytophthora foot rot, cold, calcareous soil, this rootstock grows well in sandy and heavy soils as well as its high tolerance to chlorosis and salinity. The trees budded on Cleopatra mandarin are slowly grown

until they reach the fruiting stage. Citrus seeds of all rootstock types germinate unevenly, which resulting poor seedling uniformity. Thus, it is highly interested by nursery owners to get higher and earlier germination (Llosa *et al.*, 2009; Anjum, 2010; Lacey, 2012; Sharaf *et al.*, 2016 and Al-Janabi, 2018).

GA₃ play important roles in many essential plant growth and development processes including seeds germination, stem elongation, and leaf expansion. They are often used to overcome seeds dormancy and can significantly improve seeds germination in many species, mainly through the activation of embryo growth, mobilization of reserves, and weakening of the endosperm layer. It has also been reported that seeds priming with GA₃ improves germination and the growth parameters of shoot length, root length, and seedling weight. (Khan and Chaudhry, 2006; Baskin and Baskin, 2014; Pallaoro *et al.*, 2016; Ma *et al.*, 2018 and Cipcigan *et al.*, 2020)

This study aimed to enhance the Cleopatra mandarin seeds germination process, as well as seedlings vigor by using GA₃ with different concentrations.

MATERIAL AND METHODS

The present investigation has been carried out throughout two successive seasons (2020 and 2021) on Cleopatra mandarin seeds (*Citrus reshni* L.) to enhance the Cleopatra mandarin seeds germination process, as well as seedlings vigor by seeds soaking for 24 hours in GA₃ with different concentrations (2000, 2500 and 3000 ppm) and soaking in tap water (control treatment). Also, before subjecting seeds to the differential pre-sowing treatments, they were soaked for 24 hours in Carpdazim fungicide solution at 50% concentration.

At late February 2020 and 2021 years seeds of Cleopatra mandarin rootstock were separately subjected to the pre-sowing treatments just before planting in germination boxes during 1st & 2nd experimental seasons, respectively. The seeds planted in an individual germination box previously filled with planting medium consisting of sand + clay mixture at equal proportion (v:v).

The experimental seeds were sowing in greenhouse at private orchard at Nobaria region, El Behera Governorate Egypt. The above treatments were applied as soaking in a solution of gibberellic acid in different concentrations. All the seeds of this study received the same horticultural practices except experimental treatments.

The tested treatments were evaluated through the following parameters:

Germination parameters

During each experimental season 10 seeds germinated measurements i.e., number and percentage of germinated seeds per every replicate (an individual germination box) were periodically counted at 30 and 60 days. Then average value of either number or percentage of germinated seeds per each investigated pre-sowing soak treatment was estimated as an average of its five replicates during every experimental season.

Average plant height (cm) and leaves number

On August during 2020 and 2021 years, respectively the average seedling height in cm.

was counted for 10 seedlings per each replicate (an individual germination box), then an average plant height per every investigated pre-sowing soak treatment was estimated (as an average of its five replicates). Also, leaves number were counted.

Leaf area (cm²)

On August during 2020 and 2021 years, respectively, the leaf area (cm²) was determined using discs of the leaf blades and total leaves area according to **Bremner and Taha (1966)**.

Leaf total chlorophyll

On August during 2020 and 2021 years, respectively, the leaf total chlorophyll was recorded in fresh leaves per each seedling using a portable chlorophyll meter SPAD 502 according to **Yadava, (1986)**.

Leaf chemical composition

On August during 2020 and 2021 years, respectively, five dried leaves were finely grinded and digested using micro-keildahl unit. The percentage of nitrogen content was determined according to **Naguib (1969)**. Phosphorus percentage was determined according to **AOAC, 1985**. Potassium percentage was determined according to **Brown and Lillil (1964)**.

Statistical analysis

The experimental design was randomized complete block design (RCBD) with five replicates and 40 seeds for each replicate. The data obtained were statistically analyzed using the analysis of variance method as reported by **Snedecor and Cochran, 1980**. The differences between means were differentiated by using Duncan's range test (**Duncan, 1958**).

RESULTS AND DISCUSSION

The obtained data in Table (1) showed that, all gibberellic acid concentration had a great positive effect on germination parameters after 30 and 60 days in both seasons.

As for in the first season, the number of germination seedling after 30 days in Table (1) reached 58.56 and 55.85 with 3000 and 2500 ppm of GA₃, respectively compared to the control, which was 28.96. In addition, germination percentage after 30 days gained 29.28 and 27.93 % with 3000 and 2500 ppm of GA₃, respectively compared to the control,

which was 14.48%. At the same time, number of Germination seedling after 60 days were 169.20 and 146.24 with 3000 and 2500 ppm of GA₃, respectively compared to the control, which was 49.17. Moreover, germination percentage after 60 days gained 84.60 and 73.12% with 3000 and 2500 ppm of GA₃, respectively compared to the control, which was 24.59%. The same trend was obtained with germination parameters at the second season.

In this respect, our results are in agreement with those obtained by other researchers (Ollitrault *et al.*, 2007; Turgutoglu *et al.*, 2015; Sharaf, *et al.*, 2016 and Dilip *et al.*, 2017).

Moreover, the mod of action for gibberellins depends on their binding to a carrier protein in

the first place, which allows it to be inserted into the cell, after binding to a specific receptor, it passes into the cell nucleus and changes the synthesis of genetic material (RNA). At the same time, the gibberellins in the seed induce expression by the α -amylase genes, an enzyme responsible for the hydrolysis of starch to produce simple sugars that will be the seedling's food source during germination until it can perform photosynthesis on its own. In addition, GA₃ can significantly improve seeds germination through activation the embryo growth, mobilization of reserves, and weakening of the endosperm layer (Pallaoro *et al.*, 2016; Ma *et al.*, 2018 and Cipcigan *et al.*, 2020).

Table 1. Impact of gibberellic acid treatments on number of germination seedling and germination percentage after 30 and 60 days from sowing of Cleopatra mandarin seeds (2020-2021 seasons)

Characteristics Treatments	Number of sowing seeds at start	Number of germination seedling after 30 days	Germination % after 30 days	Number of germination seedling after 60 days	Germination % after 60 days
First season 2020					
Control (water tap)	200	28.96 C	14.48 C	49.17 D	24.59 D
GA ₃ at 2000 ppm	200	48.46 B	24.23 B	107.30 C	53.65 C
GA ₃ at 2500 ppm	200	55.85 A	27.93 A	146.24 B	73.12 B
GA ₃ at 3000 ppm	200	58.56 A	29.28 A	169.20 A	84.60 A
Second season 2021					
Control (water tap)	200	32.41 C	16.20 C	47.63 D	23.82 D
GA ₃ at 2000 ppm	200	54.18 B	27.09 B	117.16 C	58.58 C
GA ₃ at 2500 ppm	200	63.40 A	31.70 A	149.15 B	74.57 B
GA ₃ at 3000 ppm	200	67.39 A	33.69 A	165.16 A	82.58 A

Mean followed by the same letter\ s within each column are not significantly different from each other at 0.5% level.

The present data in Table (2) illustrated that, all gibberellic acid treatments had a great statically influence on plant height, leaves number, leaf area, total leaves area and total chlorophyll characteristics in both seasons.

Regarding, plant height achieved 9.71 and 9.02 cm with 3000 and 2500 ppm of GA₃, respectively compared to the control, which was 6.15 cm in the first season. In addition, leaves number earned 8.75 and 8.33 with 3000 and 2500 ppm of GA₃, respectively compared to the

control, which was 6.37. At the same time, leaf area was 8.83 and 8.51cm² with 3000 and 2500 ppm of GA₃, respectively compared to the control, which was 7.22 cm² in the first season. Moreover, total leaves area attained to 77.32 and 70.96 cm² with 3000 and 2500 ppm of GA₃, respectively compared to the control, which was 46.04 cm² in the first season. The total chlorophyll acquired 68.48 and 67.28 with 3000 and 2500 ppm of GA₃, respectively compared to the control, which was 60.44. The same trend was obtained with plant height, leaves number,

leaf area, total leaves area and total chlorophyll characteristics at the second season. These results are in congeniality with those found by **Francis and Sorrell (2001); Debaje *et al.* (2010); Mahmoud (2012); Sharaf *et al.* (2016) and Dilip *et al.* (2017).**

The existing data in Table (3) supply that, all gibberellic acid treatments had a major statically impact on N, P and K leaf composition in both seasons.

Generally, leaf nitrogen content reached to 1.771 and 1.755 % with 3000 and 2500 ppm of GA₃, respectively compared to the control, which was 1.427 % in the first season. In

addition, leaf phosphorus content gained 0.393 and 0.384 % with 3000 and 2500 ppm of GA₃, respectively compared to the control, which was 0.324 %. At the same time, leaf potassium content was 2.171 and 2.130% with 3000 and 2500 ppm of GA₃, respectively compared to the control, which was 1.885 % in the first season. the same trend was obtained with N, P and K leaf composition at the second season.

These results are in general agreement with the earlier findings of **Abd El-momein *et al.*, 2007; Panigrahi and Strivastava, 2011; Mahmoud, 2012 and Sharaf *et al.*, 2016.**

Table 2. Impact of gibberellic acid treatments on plant height, leaves number, leaf area, total leaves area and total chlorophyll of Cleopatra mandarin seedling (2020-2021 seasons)

Characteristics Treatments	Plant height (cm)	Leaves number	Leaf area (cm ²)	Total leaves area (cm ²)	Total chlorophyll SPAD
First season 2020					
Control (water tap)	6.15 C	6.37 C	7.22 D	46.04 D	60.44 C
GA ₃ at 2000 ppm	7.41 B	7.60 B	8.03 C	61.01 C	63.60 B
GA ₃ at 2500 ppm	9.02 A	8.33 A	8.51 B	70.96 B	67.28 A
GA ₃ at 3000 ppm	9.71 A	8.75 A	8.83 A	77.32 A	68.48 A
Second season 2021					
Control (water tap)	6.42 C	6.39 C	7.24 D	46.26 D	59.64 C
GA ₃ at 2000 ppm	8.51 B	7.50 B	8.06 C	60.46 C	63.19 B
GA ₃ at 2500 ppm	10.18 A	8.47 A	8.48 B	71.85 B	67.96 A
GA ₃ at 3000 ppm	10.60 A	8.75 A	8.73 A	76.46 A	68.80 A

Mean followed by the same letter\ within each column are not significantly different from each other at 0.5% level.

Table 3. Impact of gibberellic acid treatments on leaves nitrogen, phosphorus and potassium contents of Cleopatra mandarin leaves (2020-2021 seasons)

Characteristics Treatments	Nitrogen percentage	Phosphorus percentage	Potassium percentage
First season 2020			
Control (water tap)	1.427 C	0.324 D	1.885 D
GA ₃ at 2000 ppm	1.589 B	0.357 C	1.995 C
GA ₃ at 2500 ppm	1.755 A	0.384 B	2.130 B
GA ₃ at 3000 ppm	1.771 A	0.393 A	2.171 A
Second season 2021			
Control (water tap)	1.426 C	0.337 C	1.877 C
GA ₃ at 2000 ppm	1.565 B	0.355 B	2.000 B
GA ₃ at 2500 ppm	1.753 A	0.382 A	2.150 A
GA ₃ at 3000 ppm	1.777 A	0.391 A	2.178 A

Mean followed by the same letter\ within each column are not significantly different from each other at 0.5% level

CONCLUSION

On the basis of present investigation, it is concluded that Cleopatra mandarin seeds treated by soaking in gibberellic acid with 2500 or 3000 ppm for 24 hours results in quicker germination, vigor vegetative growth and better germination percentage of seedlings.

REFERENCES

- Abd El-momein, Eman A. A.; Abd El-migeed, M. M. and Omayma M. M. Ismail (2007).** GA₃ and zinc sprays for improving yield and fruit quality of Washington Naval orange trees grown under sandy soil conditions. *Research Journal of Agriculture and Biological Sciences*, 3 (5): 498-503.
- Al-Janabi, A. M. I. (2018).** Effect of foliar application with kinetin and urea on some growth characteristics of Cleopatra Mandarin (*Citrus reshni* Hort. ex Tan.) rootstock. *Journal of Research in Ecology*, 6 (2): 1952-1964.
- Anjum, M. A. (2010).** Response of Cleopatra Mandarin seedlings to a polyamine-biosynthesis inhibitor under salt stress. *Acta Physiologiae Plantarum*, 32(5): 951-959.
- AOAC. (1985).** Official Methods of Analysis of the Association of Official Agriculture Chemists. 13th Ed. Benjamin Franklin Station, Washington, D. C., B. O. Box450, USA.
- Baskin, C. C. and Baskin, J. M. (2014).** Seeds: Ecology, Biogeography, and Evolution of Dormancy and Germination (2nd edition). Academic Press, San Diego.
- Bremner, P. M. and Taha, M. A. (1966).** Studies in potato agronomy 1- The effects of variety, seed size and spacing on growth, development and yield. *The Journal of Agricultural Science*, 66(2): 241-252.
- Brown, J. D. and Lilliland, O. (1964).** Rapid determination of potassium and sodium in plant material and soil extracts by flame-photometry. *The American Society for Horticultural Science*, 48: 341-346.
- Cipcigan, M. C.; Doru, P.; Cristian, R. Sisea and Rodica, M. (2020).** Gibberellic acid can improve seed germination and ornamental quality of selected cyclamen species grown under short and long days. *Agronomy*, 10: 1-19.
- Debaje, P. P.; Shinde, E. D. and Ingle, H. V. (2010).** Effect of plant growth regulators on growth and yield of acid lime (*Citrus aurantifolia*. S.). *Green Farming*, 1 (1): 62-63.
- Dilip, W. S.; Singh, D.; Moharana, D.; Rout, S. and Patra, S. S. (2017).** Influence of gibberellic acid (GA₃) on seed germination and seedling growth of Kagzi lime. *Journal of Scientific Agriculture*, 1: 62-69.
- Duncan, D. B. (1958).** Multiple range and multiple "F" test. *Biometrics*, 11: 1- 42.
- Francis, D. and Sorrell, D. A. (2001).** The interface between the cell cycle and plant growth regulators: a mini review. *Plant Growth Regulation*, 33: 1-12.
- Khan, A. S. and Chaudhry, N. Y. (2006).** GA₃ improves flower yield in some cucurbits treated with lead and mercury. *African Journal of Biotechnology*, 5(2): 149–153.
- Lacey, K. (2012).** Citrus rootstocks for WA. Department of Agriculture and Food. Farmnote: 155. Available from www.agric.wa.gov.au.
- Llosa, M. J.; Bermejo, A.; Quinones, A. C. and Forner-giner, M. A. (2009).** The citrus rootstocks Cleopatra Mandarin, *Poncirus trifoliata*, forner-alcaide 5 and forner-Alcaide 13 vary in susceptibility to iron deficiency chlorosis. *Journal of the American Pomological Society*, 63(4): 160-167.
- Ma, H. Y.; Zhao, D. D.; Ning, Q. R.; Wei, J. P.; Li, Y.; Wang, M. M.; Liu, X. L.; Jiang, C. J. and Liang, Z. W. (2018).** A Multi-year beneficial effect of seed Priming with gibberellic Acid-3 (GA₃) on plant growth and production in a perennial grass, *Leymus chinensis*. *Scientific Reports*, 8: 1-9.
- Mahmoud, T. A. (2012).** Effect of some irrigation and growth regulator treatments on growth and fruiting of Washington navel orange trees. Ph. D. thesis, Faculty of Agriculture, Zagazig University, Egypt, 208.
- Naguib, M. L. (1969).** Colorimetric determination of nitrogen components of plant tissues. *Bulletin of the Faculty of Agriculture Science, Cairo University*, 43:1.
- Ollitrault, P.; Guo, W. and Grosser, J. W. (2007).** Somatic hybridization, p. 235–260. In: I. Khan (ed.). *Citrus genetics, breeding and biotechnology*. CABI, New York.
- Pallaoro, D. S.; Avelino, A. C.; Camili, E. C. and Guimaraes, S. (2016).** Priming corn seeds

with plant growth regulator. *Journal of Seed Science*, 38: 227–232.

Panigrahi, P. and Strivastava, A. K. (2011). Deficit irrigation (DI) scheduling for mature Nagpur mandarin (*Citrus reticulata* Blanco) trees of central India. *Indian Journal of Soil Conservation*, 39(2): 149-154.

Sharaf, M. M.; Atawia, A. R.; Bakry, K. A. and EL-Rouby, M. Z. (2016). Effect of pre-sowing seeds soak in different GA₃ and ZnSo₄ solutions on germination and growth of Cleopatra Mandarin and Rangpur Lime rootstocks. *Middle East Journal of Agriculture Research*, 5 (2): 233-238.

Snedecor, G.W. and Cochran, W. G. (1980). *Statistical Methods*. Oxford and J. B. H. Publishing Co. 7th Ed. Iowa State University, Press, Am., Iowa, USA.

Turgutoglu, E.; Kurt, S. and Demir, G. (2015). Effect of GA₃ concentrations in basal medium on embryos germination of Cleopatra mandarin x carrizo citrange and Cleopatra mandarin x Flying Dragon. *Journal of Crop Breeding and Genetics*, 1(1): 17-19.

Yadava, U. L. (1986). Rapid non-destructive methods to determine chlorophyll in intact leaves. *Horticultural Science*, 21: 1449-1450.